

Supplementary Materials

Genetic identification and reiterated captures suggest that the *Astyanax mexicanus* El Pachón cavefish population is closed and declining

Laurent Legendre^{1,#}, Julie Rode^{2,#}, Isabelle Germon^{1,#}, Marie Pavie^{3,#}, Carla Quiviger¹, Maxime Pollicarpo^{1,\$}, Julien Leclercq³, Stéphane Père³, Julien Fumey^{4,5}, Carole Hyacinthe⁶, Patricia Ornelas-García⁷, Luis Espinasa⁸, Sylvie Rétaux^{3,*}, Didier Casane^{1,9,*}

¹ Université Paris-Saclay, CNRS, IRD, UMR Évolution, Génomes, Comportement et Écologie, Gif-sur-Yvette 91190, France

² Objectif Sciences International NGO, OSI-Panthera Program, Geneva, Switzerland

³ Institut de Neuroscience Paris-Saclay, Université Paris-Saclay and CNRS, Saclay 91400, France

⁴ Institut Pasteur, Université Paris Cité, CNRS UMR 2000, Microbial Paleogenomics Unit, Paris 75015, France

⁵ Institut Pasteur, Université Paris Cité, Bioinformatics and Biostatistics Hub, Paris F-75015, France

⁶ Department of Genetics, Harvard Medical School, Blavatnik Institute, Boston MA 02115, USA

⁷ Colección Nacional de Peces, Departamento de Zoología, Instituto de Biología, Universidad Nacional Autónoma de México, México DF CP 04510, México

⁸ School of Science, Marist College, Poughkeepsie, New York 12601, USA

⁹ Université Paris Cité, UFR Sciences du Vivant, Paris 75013, France


#Authors contributed equally to this work

\$Current affiliation: Zoological Institute, Department of Environmental Sciences, University of Basel, Basel, Switzerland


*Corresponding authors, E-mail: sylvie.retaux@cnrs.fr; didier.casane@universite-paris-saclay.fr

Supplementary Figure S1. Analysis of allele frequency differences between sampling sessions using Fisher's exact tests. A) without correction for multiple tests, B) with Bonferroni correction, C) with Benjamini-Hochberg procedure to decrease the false discovery rate.


A)




B)




C)




Supplementary Figure S2. Distribution of pairwise genetic distances (number of allelic differences). A) El Pachón 2016; B) El Pachón 2019; C) El Pachón 22 February 2022; D) El Pachón 25 February 2022; E) Arroyo Tampemole 2016; F) Pozo Pachón Praxedis Guerrero 2016; G) Expected distribution of pairwise genetic distances according to genealogical relationship in El Pachón 2016; H) Expected distribution of pairwise genetic distances according to genealogical relationship in El Pachón 2019; I) Expected distribution of pairwise genetic distances according to genealogical relationship in El Pachón 22 February 2022; J) Expected distribution of pairwise genetic distances according to genealogical relationship in El Pachón 25 February 2022; K) Expected distribution of pairwise genetic distances according to genealogical relationship in Arroyo Tampemole 2016; L) Expected distribution of pairwise genetic distances according to genealogical relationship in Pozo Pachón Praxedis Guerrero 2016.




G) Pachón 2016




H) Pachón 2019




I) Pachón 22 February 2022




J) Pachón 25 February 2022



K) Arroyo Tampe mole 2016



L) Pozo Pachón Praxedis Guerrero 2016



Supplementary Table S1: Sequences of the primers used to amplify the 18 microsatellite loci.

Supplementary Table S2: Fin clips and swabs genotypes obtained in the present study. Genotypes highlighted

in red are incomplete and were not used in demographic analyses.

Supplementary Table S3: Allele frequencies in A) Arroyo Tampemole 2016 and B) Pozo Pachón Praxedis Guerrero 2016.

Supplementary Table S4: Allele lengths and frequencies in El Pachón cave in 2008 as reported in Bradic *et al.* 2012.

Supplementary Table S5: Matrix of genetic distances between El Pachón genotypes obtained in 2016, 2019 and 2022.

Supplementary Table S6: Reliability of inferences of different kinships using ML Relate and according to allelic frequencies estimated during different sampling sessions.

Supplementary Table S1–S6 are listed as separate Excel files due to their large size.